



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

BEHAVIORAL HEALTH DESCRIPTION	ICD 10 CODE
Acrophobia	F40.241
Acute stress reaction	F43.0
Adjustment disorder with anxiety	F43.22
Adjustment disorder with depressed mood	F43.21
Adjustment disorder with disturbance of conduct	F43.24
Adjustment disorder with mixed anxiety and depressed mood	F43.23
Adjustment disorder with mixed disturbance of emotions and conduct	F43.25
Adjustment disorder with other symptoms	F43.29
Adjustment disorder, unspecified	F43.20
Adjustment insomnia	F51.02
Adult onset fluency disorder	F98.5
Agoraphobia with panic disorder	F40.01
Agoraphobia without panic disorder	F40.02
Agoraphobia, unspecified	F40.00
Alcohol abuse with alcohol-induced anxiety disorder	F10.180
Alcohol abuse with alcohol-induced mood disorder	F10.14
Alcohol abuse with alcohol-induced psychotic disorder with delusions	F10.150
Alcohol abuse with alcohol-induced psychotic disorder with hallucinations	F10.151
Alcohol abuse with alcohol-induced psychotic disorder, unspecified	F10.159
Alcohol abuse with alcohol-induced sexual dysfunction	F10.181
Alcohol abuse with alcohol-induced sleep disorder	F10.182
Alcohol abuse with intoxication delirium	F10.121
Alcohol abuse with other alcohol-induced disorder	F10.188
Alcohol abuse with unspecified alcohol-induced disorder	F10.19
Alcohol dependence with alcohol-induced anxiety disorder	F10.280
Alcohol dependence with alcohol-induced mood disorder	F10.24
Alcohol dependence with alcohol-induced persisting amnesic disorder	F10.26
Alcohol dependence with alcohol-induced persisting dementia	F10.27
Alcohol dependence with alcohol-induced psychotic disorder with delusions	F10.250



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Alcohol dependence with alcohol-induced psychotic disorder with hallucinations	F10.251
Alcohol dependence with alcohol-induced psychotic disorder, unspecified	F10.259
Alcohol dependence with alcohol-induced sexual dysfunction	F10.281
Alcohol dependence with alcohol-induced sleep disorder	F10.282
Alcohol dependence with intoxication delirium	F10.221
Alcohol dependence with other alcohol-induced disorder	F10.288
Alcohol dependence with unspecified alcohol-induced disorder	F10.29
Alcohol dependence with withdrawal delirium	F10.231
Alcohol dependence with withdrawal with perceptual disturbance	F10.232
Alcohol dependence with withdrawal, unspecified	F10.239
Alcohol use, unspecified with alcohol-induced anxiety disorder	F10.980
Alcohol use, unspecified with alcohol-induced mood disorder	F10.94
Alcohol use, unspecified with alcohol-induced persisting amnestic disorder	F10.96
Alcohol use, unspecified with alcohol-induced persisting dementia	F10.97
Alcohol use, unspecified with alcohol-induced psychotic disorder with delusions	F10.950
Alcohol use, unspecified with alcohol-induced psychotic disorder with hallucinations	F10.951
Alcohol use, unspecified with alcohol-induced psychotic disorder, unspecified	F10.959
Alcohol use, unspecified with alcohol-induced sexual dysfunction	F10.981
Alcohol use, unspecified with alcohol-induced sleep disorder	F10.982
Alcohol use, unspecified with intoxication delirium	F10.921
Alcohol use, unspecified with other alcohol-induced disorder	F10.988
Amnestic disorder due to known physiological condition	F04
Androphobia	F40.290
Anorexia nervosa, binge eating/purging type	F50.02
Anorexia nervosa, restricting type	F50.01
Anorexia nervosa, unspecified	F50.00
Antisocial personality disorder	F60.2
Anxiety disorder due to known physiological condition	F06.4



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Anxiety disorder, unspecified	F41.9
Arachnophobia	F40.210
Asperger's syndrome	F84.5
Attention-deficit hyperactivity disorder, combined type	F90.2
Attention-deficit hyperactivity disorder, other type	F90.8
Attention-deficit hyperactivity disorder, predominantly hyperactive type	F90.1
Attention-deficit hyperactivity disorder, predominantly inattentive type	F90.0
Attention-deficit hyperactivity disorder, unspecified type	F90.9
Autistic disorder	F84.0
Avoidant personality disorder	F60.6
Avoidant/restrictive food intake disorder	F50.82
Binge eating disorder	F50.81
Bipolar disorder, current episode depressed, mild	F31.31
Bipolar disorder, current episode depressed, mild or moderate severity, unspecified	F31.30
Bipolar disorder, current episode depressed, moderate	F31.32
Bipolar disorder, current episode depressed, severe, with psychotic features	F31.5
Bipolar disorder, current episode depressed, severe, without psychotic features	F31.4
Bipolar disorder, current episode hypomanic	F31.0
Bipolar disorder, current episode manic severe with psychotic features	F31.2
Bipolar disorder, current episode manic without psychotic features, mild	F31.11
Bipolar disorder, current episode manic without psychotic features, moderate	F31.12
Bipolar disorder, current episode manic without psychotic features, severe	F31.13
Bipolar disorder, current episode manic without psychotic features, unspecified	F31.10
Bipolar disorder, current episode mixed, mild	F31.61
Bipolar disorder, current episode mixed, moderate	F31.62
Bipolar disorder, current episode mixed, severe, with psychotic features	F31.64



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Bipolar disorder, current episode mixed, severe, without psychotic features	F31.63
Bipolar disorder, current episode mixed, unspecified	F31.60
Bipolar disorder, currently in remission, most recent episode unspecified	F31.70
Bipolar disorder, in full remission, most recent episode depressed	F31.76
Bipolar disorder, in full remission, most recent episode hypomanic	F31.72
Bipolar disorder, in full remission, most recent episode manic	F31.74
Bipolar disorder, in full remission, most recent episode mixed	F31.78
Bipolar disorder, in partial remission, most recent episode depressed	F31.75
Bipolar disorder, in partial remission, most recent episode hypomanic	F31.71
Bipolar disorder, in partial remission, most recent episode manic	F31.73
Bipolar disorder, in partial remission, most recent episode mixed	F31.77
Bipolar disorder, unspecified	F31.9
Bipolar II disorder	F31.81
Body dysmorphic disorder	F45.22
Borderline personality disorder	F60.3
Brief psychotic disorder	F23
Bulimia nervosa	F50.2
Cannabis abuse with cannabis-induced anxiety disorder	F12.180
Cannabis abuse with intoxication delirium	F12.121
Cannabis abuse with intoxication with perceptual disturbance	F12.122
Cannabis abuse with other cannabis-induced disorder	F12.188
Cannabis abuse with psychotic disorder with delusions	F12.150
Cannabis abuse with psychotic disorder with hallucinations	F12.151
Cannabis abuse with psychotic disorder, unspecified	F12.159
Cannabis abuse with unspecified cannabis-induced disorder	F12.19
Cannabis dependence with cannabis-induced anxiety disorder	F12.280
Cannabis dependence with intoxication delirium	F12.221
Cannabis dependence with intoxication with perceptual disturbance	F12.222
Cannabis dependence with other cannabis-induced disorder	F12.288
Cannabis dependence with psychotic disorder with delusions	F12.250
Cannabis dependence with psychotic disorder with hallucinations	F12.251
Cannabis dependence with psychotic disorder, unspecified	F12.259



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Cannabis dependence with unspecified cannabis-induced disorder	F12.29
Cannabis use, unspecified with anxiety disorder	F12.980
Cannabis use, unspecified with intoxication delirium	F12.921
Cannabis use, unspecified with intoxication with perceptual disturbance	F12.922
Cannabis use, unspecified with other cannabis-induced disorder	F12.988
Cannabis use, unspecified with psychotic disorder with delusions	F12.950
Cannabis use, unspecified with psychotic disorder with hallucinations	F12.951
Cannabis use, unspecified with psychotic disorder, unspecified	F12.959
Catatonic disorder due to known physiological condition	F06.1
Catatonic schizophrenia	F20.2
Childhood emotional disorder, unspecified	F93.9
Childhood onset fluency disorder	F80.81
Chronic motor or vocal tic disorder	F95.1
Claustrophobia	F40.240
Cocaine abuse with cocaine-induced anxiety disorder	F14.180
Cocaine abuse with cocaine-induced mood disorder	F14.14
Cocaine abuse with cocaine-induced psychotic disorder with delusions	F14.150
Cocaine abuse with cocaine-induced psychotic disorder with hallucinations	F14.151
Cocaine abuse with cocaine-induced psychotic disorder, unspecified	F14.159
Cocaine abuse with cocaine-induced sexual dysfunction	F14.181
Cocaine abuse with cocaine-induced sleep disorder	F14.182
Cocaine abuse with intoxication with delirium	F14.121
Cocaine abuse with intoxication with perceptual disturbance	F14.122
Cocaine abuse with other cocaine-induced disorder	F14.188
Cocaine abuse with unspecified cocaine-induced disorder	F14.19
Cocaine dependence with cocaine-induced anxiety disorder	F14.280
Cocaine dependence with cocaine-induced mood disorder	F14.24
Cocaine dependence with cocaine-induced psychotic disorder with delusions	F14.250
Cocaine dependence with cocaine-induced psychotic disorder with hallucinations	F14.251



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Cocaine dependence with cocaine-induced psychotic disorder, unspecified	F14.259
Cocaine dependence with cocaine-induced sexual dysfunction	F14.281
Cocaine dependence with cocaine-induced sleep disorder	F14.282
Cocaine dependence with intoxication delirium	F14.221
Cocaine dependence with intoxication with perceptual disturbance	F14.222
Cocaine dependence with other cocaine-induced disorder	F14.288
Cocaine dependence with unspecified cocaine-induced disorder	F14.29
Cocaine use, unspecified with cocaine-induced anxiety disorder	F14.980
Cocaine use, unspecified with cocaine-induced mood disorder	F14.94
Cocaine use, unspecified with cocaine-induced psychotic disorder with delusions	F14.950
Cocaine use, unspecified with cocaine-induced psychotic disorder with hallucinations	F14.951
Cocaine use, unspecified with cocaine-induced psychotic disorder, unspecified	F14.959
Cocaine use, unspecified with cocaine-induced sexual dysfunction	F14.981
Cocaine use, unspecified with cocaine-induced sleep disorder	F14.982
Cocaine use, unspecified with intoxication delirium	F14.921
Cocaine use, unspecified with intoxication with perceptual disturbance	F14.922
Cocaine use, unspecified with other cocaine-induced disorder	F14.988
Conduct disorder confined to family context	F91.0
Conduct disorder, adolescent-onset type	F91.2
Conduct disorder, childhood-onset type	F91.1
Conduct disorder, unspecified	F91.9
Conversion disorder with mixed symptom presentation	F44.7
Conversion disorder with motor symptom or deficit	F44.4
Conversion disorder with seizures or convulsions	F44.5
Conversion disorder with sensory symptom or deficit	F44.6
Cyclothymic disorder	F34.0
Delirium due to known physiological condition	F05
Delusional disorders	F22



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Dementia in other diseases classified elsewhere with behavioral disturbance	F02.81
Dementia in other diseases classified elsewhere without behavioral disturbance	F02.80
Dependent personality disorder	F60.7
Depersonalization-derealization syndrome	F48.1
Disinhibited attachment disorder of childhood	F94.2
Disorganized schizophrenia	F20.1
Disruptive mood dysregulation disorder	F34.81
Dissociative amnesia	F44.0
Dissociative and conversion disorder, unspecified	F44.9
Dissociative fugue	F44.1
Dissociative identity disorder	F44.81
Dissociative stupor	F44.2
Dyspareunia not due to a substance or known physiological condition	F52.6
Dysthymic disorder	F34.1
Eating disorder, unspecified	F50.9
Encopresis not due to a substance or known physiological condition	F98.1
Enuresis not due to a substance or known physiological condition	F98.0
Excoriation (skin-picking) disorder	F42.4
Exhibitionism	F65.2
Factitious disorder with combined psychological and physical signs and symptoms	F68.13
Factitious disorder with predominantly physical signs and symptoms	F68.12
Factitious disorder with predominantly psychological signs and symptoms	F68.11
Factitious disorder, unspecified	F68.10
Fear of blood	F40.230
Fear of bridges	F40.242
Fear of flying	F40.243
Fear of injections and transfusions	F40.231
Fear of injury	F40.233
Fear of other medical care	F40.232
Fear of thunderstorms	F40.220



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Female orgasmic disorder	F52.31
Female sexual arousal disorder	F52.22
Fetishism	F65.0
Frotteurism	F65.81
Gender identity disorder in adolescence and adulthood	F64.1
Gender identity disorder of childhood	F64.2
Gender identity disorder, unspecified	F64.9
Generalized anxiety disorder	F41.1
Gynephobia	F40.291
Hallucinogen abuse with hallucinogen persisting perception disorder (flashbacks)	F16.183
Hallucinogen abuse with hallucinogen-induced anxiety disorder	F16.180
Hallucinogen abuse with hallucinogen-induced mood disorder	F16.14
Hallucinogen abuse with hallucinogen-induced psychotic disorder with delusions	F16.150
Hallucinogen abuse with hallucinogen-induced psychotic disorder with hallucinations	F16.151
Hallucinogen abuse with hallucinogen-induced psychotic disorder, unspecified	F16.159
Hallucinogen abuse with intoxication with delirium	F16.121
Hallucinogen abuse with intoxication with perceptual disturbance	F16.122
Hallucinogen abuse with other hallucinogen-induced disorder	F16.188
Hallucinogen abuse with unspecified hallucinogen-induced disorder	F16.19
Hallucinogen dependence with hallucinogen persisting perception disorder (flashbacks)	F16.283
Hallucinogen dependence with hallucinogen-induced anxiety disorder	F16.280
Hallucinogen dependence with hallucinogen-induced mood disorder	F16.24
Hallucinogen dependence with hallucinogen-induced psychotic disorder with delusions	F16.250
Hallucinogen dependence with hallucinogen-induced psychotic disorder with hallucinations	F16.251



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Hallucinogen dependence with hallucinogen-induced psychotic disorder, unspecified	F16.259
Hallucinogen dependence with intoxication with delirium	F16.221
Hallucinogen dependence with other hallucinogen-induced disorder	F16.288
Hallucinogen dependence with unspecified hallucinogen-induced disorder	F16.29
Hallucinogen use, unspecified with hallucinogen persisting perception disorder (flashbacks)	F16.983
Hallucinogen use, unspecified with hallucinogen-induced anxiety disorder	F16.980
Hallucinogen use, unspecified with hallucinogen-induced mood disorder	F16.94
Hallucinogen use, unspecified with hallucinogen-induced psychotic disorder with delusions	F16.950
Hallucinogen use, unspecified with hallucinogen-induced psychotic disorder with hallucinations	F16.951
Hallucinogen use, unspecified with hallucinogen-induced psychotic disorder, unspecified	F16.959
Hallucinogen use, unspecified with intoxication with delirium	F16.921
Hallucinogen use, unspecified with other hallucinogen-induced disorder	F16.988
Histrionic personality disorder	F60.4
Hoarding disorder	F42.3
Hypoactive sexual desire disorder	F52.0
Hypochondriacal disorder, unspecified	F45.20
Hypochondriasis	F45.21
Impulse disorder, unspecified	F63.9
Inhalant abuse with inhalant-induced anxiety disorder	F18.180
Inhalant abuse with inhalant-induced dementia	F18.17
Inhalant abuse with inhalant-induced mood disorder	F18.14
Inhalant abuse with inhalant-induced psychotic disorder with delusions	F18.150
Inhalant abuse with inhalant-induced psychotic disorder with hallucinations	F18.151



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Inhalant abuse with inhalant-induced psychotic disorder, unspecified	F18.159
Inhalant abuse with intoxication delirium	F18.121
Inhalant abuse with other inhalant-induced disorder	F18.188
Inhalant abuse with unspecified inhalant-induced disorder	F18.19
Inhalant dependence with inhalant-induced anxiety disorder	F18.280
Inhalant dependence with inhalant-induced dementia	F18.27
Inhalant dependence with inhalant-induced mood disorder	F18.24
Inhalant dependence with inhalant-induced psychotic disorder with delusions	F18.250
Inhalant dependence with inhalant-induced psychotic disorder with hallucinations	F18.251
Inhalant dependence with inhalant-induced psychotic disorder, unspecified	F18.259
Inhalant dependence with intoxication delirium	F18.221
Inhalant use, unspecified with inhalant-induced anxiety disorder	F18.980
Inhalant use, unspecified with inhalant-induced mood disorder	F18.94
Inhalant use, unspecified with inhalant-induced persisting dementia	F18.97
Inhalant use, unspecified with inhalant-induced psychotic disorder with delusions	F18.950
Inhalant use, unspecified with inhalant-induced psychotic disorder with hallucinations	F18.951
Inhalant use, unspecified with inhalant-induced psychotic disorder, unspecified	F18.959
Inhalant use, unspecified with intoxication with delirium	F18.921
Insomnia due to other mental disorder	F51.05
Insufficient sleep syndrome	F51.12
Intermittent explosive disorder	F63.81
Kleptomania	F63.2
Major depressive disorder, recurrent severe without psychotic features	F33.2
Major depressive disorder, recurrent, in full remission	F33.42
Major depressive disorder, recurrent, in partial remission	F33.41
Major depressive disorder, recurrent, in remission, unspecified	F33.40
Major depressive disorder, recurrent, mild	F33.0



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Major depressive disorder, recurrent, moderate	F33.1
Major depressive disorder, recurrent, severe with psychotic symptoms	F33.3
Major depressive disorder, recurrent, unspecified	F33.9
Major depressive disorder, single episode, in full remission	F32.5
Major depressive disorder, single episode, in partial remission	F32.4
Major depressive disorder, single episode, mild	F32.0
Major depressive disorder, single episode, moderate	F32.1
Major depressive disorder, single episode, severe with psychotic features	F32.3
Major depressive disorder, single episode, severe without psychotic features	F32.2
Major depressive disorder, single episode, unspecified	F32.9
Male erectile disorder	F52.21
Male orgasmic disorder	F52.32
Manic episode in full remission	F30.4
Manic episode in partial remission	F30.3
Manic episode without psychotic symptoms, mild	F30.11
Manic episode without psychotic symptoms, moderate	F30.12
Manic episode without psychotic symptoms, unspecified	F30.10
Manic episode, severe with psychotic symptoms	F30.2
Manic episode, severe, without psychotic symptoms	F30.13
Manic episode, unspecified	F30.9
Mixed obsessional thoughts and acts	F42.2
Mood disorder due to known physiological condition with depressive features	F06.31
Mood disorder due to known physiological condition with major depressive-like episode	F06.32
Mood disorder due to known physiological condition with manic features	F06.33
Mood disorder due to known physiological condition with mixed features	F06.34
Mood disorder due to known physiological condition, unspecified	F06.30
Narcissistic personality disorder	F60.81
Nightmare disorder	F51.5
Nonpsychotic mental disorder, unspecified	F48.9



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Obsessive-compulsive disorder	F42
Obsessive-compulsive personality disorder	F60.5
Opioid abuse with intoxication delirium	F11.121
Opioid abuse with intoxication with perceptual disturbance	F11.122
Opioid abuse with opioid-induced mood disorder	F11.14
Opioid abuse with opioid-induced psychotic disorder with delusions	F11.150
Opioid abuse with opioid-induced psychotic disorder with hallucinations	F11.151
Opioid abuse with opioid-induced psychotic disorder, unspecified	F11.159
Opioid abuse with opioid-induced sexual dysfunction	F11.181
Opioid abuse with opioid-induced sleep disorder	F11.182
Opioid abuse with other opioid-induced disorder	F11.188
Opioid abuse with unspecified opioid-induced disorder	F11.19
Opioid dependence with intoxication delirium	F11.221
Opioid dependence with opioid-induced mood disorder	F11.24
Opioid dependence with opioid-induced psychotic disorder with delusions	F11.250
Opioid dependence with opioid-induced psychotic disorder with hallucinations	F11.251
Opioid dependence with opioid-induced psychotic disorder, unspecified	F11.259
Opioid dependence with opioid-induced sexual dysfunction	F11.281
Opioid dependence with opioid-induced sleep disorder	F11.282
Opioid dependence with other opioid-induced disorder	F11.288
Opioid dependence with unspecified opioid-induced disorder	F11.29
Opioid use, unspecified with intoxication delirium	F11.921
Opioid use, unspecified with intoxication with perceptual disturbance	F11.922
Opioid use, unspecified with opioid-induced mood disorder	F11.94
Opioid use, unspecified with opioid-induced psychotic disorder with delusions	F11.950
Opioid use, unspecified with opioid-induced psychotic disorder with hallucinations	F11.951
Opioid use, unspecified with opioid-induced psychotic disorder, unspecified	F11.959
Opioid use, unspecified with opioid-induced sexual dysfunction	F11.981



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Opioid use, unspecified with opioid-induced sleep disorder	F11.982
Opioid use, unspecified with other opioid-induced disorder	F11.988
Oppositional defiant disorder	F91.3
Other animal type phobia	F40.218
Other bipolar disorder	F31.89
Other childhood disintegrative disorder	F84.3
Other childhood disorders of social functioning	F94.8
Other childhood emotional disorders	F93.8
Other conduct disorders	F91.8
Other depressive episodes	F32.8
Other developmental disorders of speech and language	F80.89
Other disorders of psychological development	F88
Other eating disorders	F50.8
Other feeding disorders of infancy and early childhood	F98.29
Other gender identity disorders	F64.8
Other hypochondriacal disorders	F45.29
Other impulse disorders	F63.8
Other insomnia not due to a substance or known physiological condition	F51.09
Other manic episodes	F30.8
Other mixed anxiety disorders	F41.3
Other natural environment type phobia	F40.228
Other obsessive compulsive disorder	F42.8
Other paraphilias	F65.89
Other persistent mood «affective» disorders	F34.8
Other personality and behavioral disorders due to known physiological condition	F07.89
Other pervasive developmental disorders	F84.8
Other phobic anxiety disorders	F40.8
Other psychoactive substance abuse with intoxication delirium	F19.121
Other psychoactive substance abuse with intoxication with perceptual disturbances	F19.122



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other psychoactive substance abuse with other psychoactive substance-induced disorder	F19.188
Other psychoactive substance abuse with psychoactive substance-induced mood disorder	F19.14
Other psychoactive substance abuse with psychoactive substance-induced sexual dysfunction	F19.181
Other psychoactive substance abuse with psychoactive substance-induced sleep disorder	F19.182
Other psychoactive substance abuse with psychoactive substance-induced anxiety disorder	F19.180
Other psychoactive substance abuse with psychoactive substance-induced persisting amnesic disorder	F19.16
Other psychoactive substance abuse with psychoactive substance-induced persisting dementia	F19.17
Other psychoactive substance abuse with psychoactive substance-induced psychotic disorder with delus	F19.150
Other psychoactive substance abuse with psychoactive substance-induced psychotic disorder with hallu	F19.151
Other psychoactive substance abuse with psychoactive substance-induced psychotic disorder, unspecifi	F19.159
Other psychoactive substance abuse with unspecified psychoactive substance-induced disorder	F19.19
Other psychoactive substance dependence with intoxication delirium	F19.221
Other psychoactive substance dependence with intoxication with perceptual disturbance	F19.222
Other psychoactive substance dependence with other psychoactive substance-induced disorder	F19.288



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other psychoactive substance dependence with psychoactive substance-induced anxiety disorder	F19.280
Other psychoactive substance dependence with psychoactive substance-induced mood disorder	F19.24
Other psychoactive substance dependence with psychoactive substance-induced persisting amnestic diso	F19.26
Other psychoactive substance dependence with psychoactive substance-induced persisting dementia	F19.27
Other psychoactive substance dependence with psychoactive substance-induced psychotic disorder with	F19.250
Other psychoactive substance dependence with psychoactive substance-induced psychotic disorder with	F19.251
Other psychoactive substance dependence with psychoactive substance-induced psychotic disorder, unsp	F19.259
Other psychoactive substance dependence with psychoactive substance-induced sexual dysfunction	F19.281
Other psychoactive substance dependence with psychoactive substance-induced sleep disorder	F19.282
Other psychoactive substance dependence with unspecified psychoactive substance-induced disorder	F19.29
Other psychoactive substance dependence with withdrawal delirium	F19.231
Other psychoactive substance dependence with withdrawal with perceptual disturbance	F19.232
Other psychoactive substance dependence with withdrawal, unspecified	F19.239



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other psychoactive substance use, unspecified with intoxication with delirium	F19.921
Other psychoactive substance use, unspecified with intoxication with perceptual disturbance	F19.922
Other psychoactive substance use, unspecified with other psychoactive substance-induced disorder	F19.988
Other psychoactive substance use, unspecified with psychoactive substance-induced anxiety disorder	F19.980
Other psychoactive substance use, unspecified with psychoactive substance-induced mood disorder	F19.94
Other psychoactive substance use, unspecified with psychoactive substance-induced persisting amnesti	F19.96
Other psychoactive substance use, unspecified with psychoactive substance-induced persisting dementi	F19.97
Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder	F19.950
Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder	F19.951
Other psychoactive substance use, unspecified with psychoactive substance-induced psychotic disorder	F19.959
Other psychoactive substance use, unspecified with psychoactive substance-induced sexual dysfunction	F19.981
Other psychoactive substance use, unspecified with psychoactive substance-induced sleep disorder	F19.982



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other psychoactive substance use, unspecified with withdrawal delirium	F19.931
Other psychoactive substance use, unspecified with withdrawal with perceptual disturbance	F19.932
Other psychotic disorder not due to a substance or known physiological condition	F28
Other reactions to severe stress	F43.8
Other recurrent depressive disorders	F33.8
Other schizoaffective disorders	F25.8
Other schizophrenia	F20.89
Other sexual disorders	F66
Other sexual dysfunction not due to a substance or known physiological condition	F52.8
Other situational type phobia	F40.248
Other sleep disorders not due to a substance or known physiological condition	F51.8
Other somatoform disorders	F45.8
Other specific personality disorders	F60.89
Other specified anxiety disorders	F41.8
Other specified behavioral and emotional disorders with onset usually occurring in childhood and ado	F98.8
Other specified depressive episodes	F32.89
Other specified disorders of adult personality and behavior	F68.8
Other specified eating disorder	F50.89
Other specified mental disorders due to known physiological condition	F06.8
Other specified persistent mood disorders	F34.89
Other specified phobia	F40.298
Other stimulant abuse with intoxication delirium	F15.121
Other stimulant abuse with intoxication with perceptual disturbance	F15.122
Other stimulant abuse with other stimulant-induced disorder	F15.188
Other stimulant abuse with stimulant-induced anxiety disorder	F15.180
Other stimulant abuse with stimulant-induced mood disorder	F15.14



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other stimulant abuse with stimulant-induced psychotic disorder with delusions	F15.150
Other stimulant abuse with stimulant-induced psychotic disorder with hallucinations	F15.151
Other stimulant abuse with stimulant-induced psychotic disorder, unspecified	F15.159
Other stimulant abuse with stimulant-induced sexual dysfunction	F15.181
Other stimulant abuse with stimulant-induced sleep disorder	F15.182
Other stimulant abuse with unspecified stimulant-induced disorder	F15.19
Other stimulant dependence with intoxication delirium	F15.221
Other stimulant dependence with intoxication with perceptual disturbance	F15.222
Other stimulant dependence with other stimulant-induced disorder	F15.288
Other stimulant dependence with stimulant-induced anxiety disorder	F15.280
Other stimulant dependence with stimulant-induced mood disorder	F15.24
Other stimulant dependence with stimulant-induced psychotic disorder with delusions	F15.250
Other stimulant dependence with stimulant-induced psychotic disorder with hallucinations	F15.251
Other stimulant dependence with stimulant-induced psychotic disorder, unspecified	F15.259
Other stimulant dependence with stimulant-induced sexual dysfunction	F15.281
Other stimulant dependence with stimulant-induced sleep disorder	F15.282
Other stimulant dependence with unspecified stimulant-induced disorder	F15.29
Other stimulant use, unspecified with intoxication delirium	F15.921
Other stimulant use, unspecified with intoxication with perceptual disturbance	F15.922
Other stimulant use, unspecified with other stimulant-induced disorder	F15.988
Other stimulant use, unspecified with stimulant-induced anxiety disorder	F15.980
Other stimulant use, unspecified with stimulant-induced mood disorder	F15.94



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Other stimulant use, unspecified with stimulant-induced psychotic disorder with delusions	F15.950
Other stimulant use, unspecified with stimulant-induced psychotic disorder with hallucinations	F15.951
Other stimulant use, unspecified with stimulant-induced psychotic disorder, unspecified	F15.959
Other stimulant use, unspecified with stimulant-induced sexual dysfunction	F15.981
Other stimulant use, unspecified with stimulant-induced sleep disorder	F15.982
Other tic disorders	F95.8
Pain disorder exclusively related to psychological factors	F45.41
Pain disorder with related psychological factors	F45.42
Panic Disorder (episodic paroxysmal anxiety)	F41.0
Panic disorder «episodic paroxysmal anxiety» without agoraphobia	F41.0
Paradoxical insomnia	F51.03
Paranoid personality disorder	F60.0
Paranoid schizophrenia	F20.0
Paraphilia, unspecified	F65.9
Pathological gambling	F63.0
Pedophilia	F65.4
Persistent mood «affective» disorder, unspecified	F34.9
Personality change due to known physiological condition	F07.0
Personality disorder, unspecified	F60.9
Pervasive developmental disorder, unspecified	F84.9
Phobic anxiety disorder, unspecified	F40.9
Pica of infancy and childhood	F98.3
Postconcussional syndrome	F07.81
Post-traumatic stress disorder, acute	F43.11
Post-traumatic stress disorder, chronic	F43.12
Post-traumatic stress disorder, unspecified	F43.10



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Premature ejaculation	F52.4
Premenstrual dysphoric disorder	F32.81
Primary hypersomnia	F51.11
Primary insomnia	F51.01
Pseudobulbar affect	F48.2
Psychotic disorder with delusions due to known physiological condition	F06.2
Psychotic disorder with hallucinations due to known physiological condition	F06.0
Puerperal psychosis	F53
Pyromania	F63.1
Reaction to severe stress, unspecified	F43.9
Reactive attachment disorder of childhood	F94.1
Residual schizophrenia	F20.5
Rumination disorder of infancy	F98.21
Sadomasochism, unspecified	F65.50
Schizoaffective disorder, bipolar type	F25.0
Schizoaffective disorder, depressive type	F25.1
Schizoaffective disorder, unspecified	F25.9
Schizoid personality disorder	F60.1
Schizophrenia, unspecified	F20.9
Schizophreniform disorder	F20.81
Schizotypal disorder	F21
Sedative, hypnotic or anxiolytic abuse with intoxication delirium	F13.121
Sedative, hypnotic or anxiolytic abuse with other sedative, hypnotic or anxiolytic-induced disorder	F13.188
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced anxiety disorder	F13.180
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced mood disorder	F13.14



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced psychotic disor	F13.151
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced psychotic disor	F13.159
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced psychotic disor	F13.150
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced sexual dysfunct	F13.181
Sedative, hypnotic or anxiolytic abuse with sedative, hypnotic or anxiolytic-induced sleep disorder	F13.182
Sedative, hypnotic or anxiolytic abuse with unspecified sedative, hypnotic or anxiolytic-induced dis	F13.19
Sedative, hypnotic or anxiolytic dependence with intoxication delirium	F13.221
Sedative, hypnotic or anxiolytic dependence with other sedative, hypnotic or anxiolytic-induced diso	F13.288
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced anxiety di	F13.280
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced mood disor	F13.24
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced persisting	F13.27
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced persisting	F13.26



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced psychotic	F13.259
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced psychotic	F13.251
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced psychotic	F13.250
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced sexual dys	F13.281
Sedative, hypnotic or anxiolytic dependence with sedative, hypnotic or anxiolytic-induced sleep diso	F13.282
Sedative, hypnotic or anxiolytic dependence with unspecified sedative, hypnotic or anxiolytic-induce	F13.29
Sedative, hypnotic or anxiolytic dependence with withdrawal delirium	F13.231
Sedative, hypnotic or anxiolytic dependence with withdrawal with perceptual disturbance	F13.232
Sedative, hypnotic or anxiolytic dependence with withdrawal, unspecified	F13.239
Sedative, hypnotic or anxiolytic use, unspecified with intoxication delirium	F13.921
Sedative, hypnotic or anxiolytic use, unspecified with other sedative, hypnotic or anxiolytic-induce	F13.988
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced anxi	F13.980
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced mood	F13.94



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced pers	F13.97
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced pers	F13.96
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced psyc	F13.950
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced psyc	F13.959
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced psyc	F13.951
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced sexu	F13.981
Sedative, hypnotic or anxiolytic use, unspecified with sedative, hypnotic or anxiolytic-induced slee	F13.982
Sedative, hypnotic or anxiolytic use, unspecified with withdrawal delirium	F13.931
Sedative, hypnotic or anxiolytic use, unspecified with withdrawal with perceptual disturbances	F13.932
Selective mutism	F94.0
Separation anxiety disorder of childhood	F93.0
Sexual aversion disorder	F52.1
Sexual masochism	F65.51
Sexual sadism	F65.52
Shared psychotic disorder	F24
Sleep disorder not due to a substance or known physiological condition, unspecified	F51.9
Sleep terrors [night terrors]	F51.4



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Sleepwalking [somnambulism]	F51.3
Social phobia, generalized	F40.11
Social phobia, unspecified	F40.10
Social pragmatic communication disorder	F80.82
Somatization disorder	F45.0
Somatoform disorder, unspecified	F45.9
State of emotional shock and stress, unspecified	R45.7
Stereotyped movement disorders	F98.4
Tic disorder, unspecified	F95.9
Tourette's disorder	F95.2
Transient tic disorder	F95.0
Transsexualism	F64.0
Transvestic fetishism	F65.1
Trichotillomania	F63.3
Undifferentiated schizophrenia	F20.3
Undifferentiated somatoform disorder	F45.1
Unspecified behavioral and emotional disorders with onset usually occurring in childhood and adolesc	F98.9
Unspecified behavioral syndromes associated with physiological disturbances and physical factors	F59
Unspecified dementia with behavioral disturbance	F03.91
Unspecified dementia without behavioral disturbance	F03.90
Unspecified disorder of psychological development	F89
Unspecified mental disorder due to known physiological condition	F09
Unspecified mood «affective» disorder	F39
Unspecified personality and behavioral disorder due to known physiological condition	F07.9
Unspecified psychosis not due to a substance or known physiological condition	F29



ICD-10-CM CODES (BH commonly used)

These commonly used ICD-10 diagnosis codes are intended to assist physicians and other authorized ordering parties in providing correct ICD-10 codes as required by Medicare and other insurers. The codes are based on ICD-10-CM 2018, Medicare Regulations and Manuals authorized by the Centers for Medicare and Medicaid Services. The list is not complete – the ICD-10-CM book should be referenced for a complete list of codes. The physician or other authorized ordering party is responsible for providing correct codes that support the medical necessity of each test ordered for the diagnosis and treatment of the individual patient.

Unspecified sexual dysfunction not due to a substance or known physiological condition	F52.9
Vaginismus not due to a substance or known physiological condition	F52.5
Vascular dementia with behavioral disturbance	F01.51
Vascular dementia without behavioral disturbance	F01.50
Voyeurism	F65.3